

Preparing for Your Colonoscopy (Extended CoLyte)

About the Procedure: You are having a test called a colonoscopy. This is an examination of your entire large bowel. To ensure the colon is as clean as possible, it is very important to follow the steps in the preparation instructions as it is possible to miss growths if stool is present. Your healthcare team considers your bowel empty when you are only passing liquid. The liquid may be clear, yellow or coloured with flecks of stool. If you do not complete your prep as instructed your appointment may be cancelled.

Seven (7) days before your colonoscopy

Call the doctor's office if you are taking any of the following blood-thinning medications:

Apixaban (Eliquis®)	Clopidogrel (Plavix®)
Rivaroxaban (Xarelto®)	Ticagrelor (Brilinta®)
Edoxaban (Lixiana®)	Prasugrel (Effient®)
Dabigatran (Pradaxa®)	Warfarin

- **Stop taking:** iron tablets and vitamins. Continue all other medications unless told otherwise.
- **Stop eating:** nuts and seeds, e.g., flax, sesame and chia seeds, berries and popcorn.

If you have diabetes: You should have received the *Preparing for Your Colonoscopy: Adjusting Your Diabetes Medication* handout. Please review.

Make plans to take time off work: Consider taking the day off work before the colonoscopy to complete your bowel preparation. You should plan to be off work the day of your colonoscopy.

Make transportation arrangements: You will need a trusted person to take you home after the colonoscopy procedure.

Buy two (2) containers of CoLyte (each are 4 litres) and three (3) bisacodyl (Dulcolax) tablets from the pharmacy. A prescription is not needed.

If you need clarification about your colonoscopy preparation, please contact the physician doing the procedure for direction. If your colonoscopy was booked through the Colorectal Cancer Screening Program, please call the Nurse Navigator at 1-855-733-0762.

If you have an emergency the day of the colonoscopy, you must call the Endoscopy Unit where your procedure is scheduled:

- QEH Endoscopy Unit Tel: 902-894-0055
- PCH Endoscopy Unit Tel: 902-432-2541

The day before your colonoscopy

- **Do not eat** any solid foods.
- You should **consume 3 to 4 litres of clear liquids** in addition to the bowel preparation solution to stay hydrated. You may drink the following liquids up until 2 hours before the colonoscopy:
 - Sports drinks, e.g., Electrolyte Gastro, Gatorade or Powerade
 - Light-colored soft drinks, e.g., Sprite, Ginger Ale, 7-UP
 - Water
 - Kool-Aid, iced tea, lemonade
 - Black or herbal tea, coffee without milk products or substitutes
 - Pulp-free fruit juices, e.g., apple, white grape, and white cranberry
 - Clear soups, e.g., broth/bouillon
 - Pulp-free coconut water
 - Popsicles
 - Jell-O
 - **no red, blue or purple coloured liquids; no dairy drinks; no alcohol.**

Prepare and refrigerate your bowel preparation solution according to the directions on the container. Chilling improves the taste and drinking through a straw helps bypass the taste buds.

If you have diabetes: try to maintain your regular caloric intake in liquid form. Test your blood sugar several times throughout the day including before bedtime.

What to expect after your first dose of the bowel preparation solution?

- Remain close to a toilet. You will have frequent diarrhea-type bowel movements. Cramping can occur and is considered normal.
- The solution usually starts working within 1-3 hours **but may take longer**. It may take up to 6 to 8 hours or more for some people. Your next dose will take less time to work. Keep this in mind if you are travelling the day before your procedure.
- Gravol 25-50 mg and other medications (e.g., Tylenol/migraine medication) may be taken by mouth if needed when completing the bowel preparation up until 3 hours before your colonoscopy.

Hospital Appointment and Arrival Information

Hospital Arrival

You must arrive and register in the Admitting Department 60 to 90 minutes before your scheduled colonoscopy. If your appointment is at 8:00 AM, please arrive and register at 7:00 AM.

The colonoscopy will take 20 to 30 minutes to perform, and you need to stay 30 minutes to 1 hour after it is over. Plan to be at the hospital for 2 to 3 hours from start to finish.

Hospital Checklist

- Bring a list of current medications
- Bring a translator if you do not understand or speak English
- Bring a support person if you require assistance
- Bring your PEI Health Card or proof of substitute medical insurance plan
- **Do not** wear any jewelry, e.g., rings, watches, earrings, bracelets, or necklaces
- **Do not** bring any valuables

REMINDER:

- If you received **sedation medication** for the colonoscopy, you are considered impaired for up to 24 hours after the test:
 - You should not sign any legal documents.
 - You will not be able to drive yourself home. You will need a trusted person to meet you at the Endoscopy Unit. If you do not have a driver your test will be cancelled. A taxi is acceptable only if a trusted individual is accompanying you.
 - You should not drink alcohol or take sleeping pills or anti-anxiety medications.
 - You should not be responsible for another person's care, e.g., a baby, young child, or person in poor health.
- If you take blood-thinning medication, discuss when to restart these medications after your colonoscopy with the doctor. Resume all other medications as usual unless told otherwise.

Extended CoLyte Bowel Preparation Instructions

At 2:00 PM the day before the colonoscopy:

- Take three (3) bisacodyl (Dulcolax) tablets with water, do not chew or crush the tablets. The first bowel movement usually occurs in 1 to 6 hours after swallowing the tablets.
- Ensure the two jugs of CoLyte prep solution are in the fridge and chilling.

At 4:00 PM the day before the colonoscopy:

- Drink half (2 litres) of the solution from the first jug of the CoLyte whether you have had a bowel movement or not. Drink at least 250 ml every 15 minutes. Rapid drinking of each glass is preferred. A watery bowel movement should begin in approximately one (1) hour. Be sure to drink all the solution.

At 9:00 PM the day before the colonoscopy:

- Drink the remaining solution from the first jug (2 litres) of the CoLyte. Drink at least 250 ml every 15 minutes. Rapid drinking of each glass is preferred. Be sure to drink all the solution.

Four to six (4-6) hours before the colonoscopy:

- Drink half (2 litres) of the solution from the second jug of the CoLyte. Drink at least 250 ml every 15 minutes. Rapid drinking of each glass is preferred. Be sure to drink all the solution.
Note: If you have an early morning appointment, you will need to get up in the middle of the night to finish the solution.

Two (2) hours before colonoscopy:

- No eating or drinking.